



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>A61K 31/70</b>	<b>A1</b>	(11) International Publication Number: <b>WO 97/40838</b> (43) International Publication Date: 6 November 1997 (06.11.97)
<p>(21) International Application Number: <b>PCT/SE97/00735</b></p> <p>(22) International Filing Date: 30 April 1997 (30.04.97)</p> <p>(30) Priority Data: 9601677-9      2 May 1996 (02.05.96)      SE</p> <p>(71) Applicant (for all designated States except US): <b>SCOTIA LIPIDTEKNIK AB [SE/SE]; P.O. Box 6686, S-113 84 Stockholm (SE).</b></p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): <b>VASÄNGE, Mervi [SE/SE]; Hässjövägen 20, S-752 38 Uppsala (SE). ROLFSEN, Wenche [SE/SE]; Kvarnbogatan 16, S-752 39 Uppsala (SE). BOHLIN, Lars [SE/SE]; Ekolnsvägen 13D, S-756 53 Uppsala (SE).</b></p> <p>(74) Agents: <b>LARFELDT, Helene et al.; Bergensträhle &amp; Lindvall AB, P.O. Box 17704, S-118 93 Stockholm (SE).</b></p>	<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p>(54) Title: <b>NEW USE</b></p> <p>(57) Abstract</p> <p>The invention refers to the use of a sulpholipid SQDG having formula (I), wherein R<sub>1</sub> and R<sub>2</sub>, which are the same or different, are hydrogen or saturated or unsaturated, optionally hydroxy-substituted acyl groups, protonated or as a salt, for the preparation of a pharmaceutical composition for the treatment of inflammatory skin diseases or disorders, especially psoriasis. A pharmaceutical composition comprising an SQDG compound in combination with a polyunsaturated fatty acid and a therapeutically acceptable carrier is also described.</p> <div style="text-align: center;"> <math display="block">  \begin{array}{c}  \text{CH}_2\text{OR}^1 \\    \\  \text{R}^2\text{OCH} \\    \\  \text{CH}_2\text{O} - \text{deoxyhexose} - \text{SO}_3^-  \end{array}  \quad (I)  </math> </div>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**NEW USE**

The present invention refers to a new PAF-receptor antagonist, the sulpholipid SQDG, as well as to the use of  
5 SQDG for the prophylaxis or treatment of inflammatory skin diseases, especially psoriasis.

**Background of the invention**

10 Inflammatory skin diseases such as atopic dermatitis, urticaria and especially psoriasis still constitute a great problem for the affected patients as there are today no effective therapies.

Psoriasis is a common chronic inflammatory dermatosis with a global distribution; it has been estimated that about  
15 1.5 % of the population in the western countries can be expected to suffer from the disease during their lifetime. A number of different clinical patterns of psoriasis are acknowledged, the most common being plaque psoriasis.

Hyperproliferation, inflammation with massive infiltration of leucocytes and disturbances in cell differentiation  
20 are the typical characteristics of psoriatic skin. The number of basal cells is vastly increased which reduces the turnover time for the epidermis from the normal 27 days to 3-4 days. The normal events of cell maturation and keratinization  
25 do not occur. This proliferation of keratinocytes occurs both in involved and non-involved psoriatic skin but is most pronounced in the plaques. The inflammatory infiltrate from psoriatic lesions has been found to consist predominantly of mononuclear T lymphocytes. There is a disturbed T cell  
30 function also in the circulating blood, which implies a possible cell-mediated immunological abnormality in psoriasis. The phagocytes that can be identified histologically in the psoriatic lesion, neutrophils in particular, suggest a role for the chemotactic inflammatory mediators in  
35 the pathology of psoriasis. Among these interleukin-1, -6 and -8, leukotriene B<sub>4</sub> and PAF have been isolated in pathological amounts in the affected skin.

There is a number of different therapies of psoriasis of varying effectiveness, none being perfect. Among the

antipsoriatic drugs of natural origin can be mentioned coal tar, dithranol, psoralens, retinoids and cyclosporin A. In addition to said more established therapies can also be mentioned the use of podophyllotoxin, a lignan which can be

isolated from *Podophyllum* species, polyunsaturated fatty acids, such as eicosapentaenoic acid and gamma-linolenic acid, and colchicine, an alkaloid from the crocus plant.

The use of essential fatty acids in atopic dermatitis and psoriasis, respectively, are described by Wright, S., British Journal of Dermatology 125, 503-515, 1991.

In the traditional medicine of Honduras the name Calaguala is used for the extract of a number of closely related *Polypodium* species, including *P. decumanum* Willd., *P. aureum* (or *P. leucatomos*), *P. lowei* C.Ch., *P. loriceum* L., *P. triseriale* Sw., *P. fraxinifolium* Jacq. and *P. dissimile* L (Molina, personal communication, 1991). A decoction or infusion of the Calaguala plant has been used to treat a number of diseases including peptic ulcer, kidney problems, diarrhoea and arthritis or other pains in joints and tendons.

Over the last two decades several clinical trials have been performed on Calaguala in the treatment of psoriasis, as well as atopic dermatitis (Vargas Gonzales, J.P., et al., Acta Pediatrica. Esp. 46(1), 556-561, 1988) and vitiligo (Gonzales, S., et al., J. Invest. Derm. 102(4), 651, 1994).

Platelet activating factor (PAF; 1-O-alkyl-2-acetyl-sn-glycero-3-phosphocholine) is a phospholipid derived mediator with a diversity of biological effects. PAF is released from various cell types, including platelets, neutrophils, monocytes and endothelial cells. In addition, several cell types, including neutrophils, are known to express specific PAF receptors on their cell membrane. In vitro PAF exhibits effects, including aggregation and degranulation of leukocytes and inhibition of lymphocyte proliferation. In vivo effects include hypotension, acute renal failure and increase in vascular permeability. PAF is thought to be associated with a number of pathological conditions, such as shock, airway hypersensitivity and asthma, inflammation of many different types, psoriasis, arthritis and graft rejection and various cardiovascular disorders especially

ones associated with thrombosis, blood coagulation and platelet aggregation.

A large number of natural and synthetic PAF receptor antagonists has been discovered which show different physiological effects, but to date they have been of limited usefulness for pharmaceutical purposes. The use of some compounds in the treatment of asthma is, however, currently being evaluated. As the PAF antagonist therapy is relatively new, more research is still needed for the judgement of therapeutic benefit for specific indications. It is believed that safe and potent compounds will have a substantial role in therapeutical treatments.

Perhaps the best known of all PAF antagonists is the ginkgolide BN 52021 which was isolated from the "fossil tree" *Ginkgo biloba* in the late 60's. It is a specific PAF receptor antagonist and has shown anti-inflammatory properties in several *in vivo* and *in vitro* models. For the potential oral treatment of chronic inflammatory diseases, the synthetic hexazepine derivative BN 50730 was later developed. It shows a several ten-folds more potent PAF antagonistic activity *in vitro*. (Guinot, P. in *Clinical Reviews in Allergy* 12, 397-417, 1994.)

It has been suggested that PAF contributes to the inflammatory aspects of psoriasis (Cunningham, 1990). Factors that implicate its role in the pathogenesis and symptoms are that PAF has been isolated in elevated amounts on psoriatic skin, that injection of PAF causes vasodilation and increased vascular permeability, and that PAF is chemoattractant mainly to neutrophils.

Recently, PAF activity has been measured in blood plasma of patients with psoriasis using a radioimmunoassay technique (Izaki et al., *British Journal of Dermatology* 136, 1060-1064, 1996). It was shown that levels of PAF were significantly elevated in patients with psoriasis and that when these patients were treated with conventional anti-psoriatic methods the levels were decreased. Also, the decrease was correlated with a clinical improvement. It is concluded that PAF has a role in the acute phase of psoriatic and leucotactic inflammation.

In another study, chemotactic responsiveness of peripheral blood eosinophils from healthy subjects and from patients with inflammatory dermatoses (psoriasis, atopic dermatitis) was determined. It was found that chemotactic responsiveness was significantly enhanced in severely affected patients and that this increase was not related to a specific disease. The increased responsiveness in peripheral eosinophils to PAF is suggested to be related to altered receptor expression during cutaneous inflammation (Morita et al. 1989).


#### Prior art

The sulphoquinovosyldiacylglycerols, SQDG, constitute one of the four major classes of polar lipids in chloroplast membranes of most photosynthetic organisms. They are structural components of the thylakoid membrane and are involved in the electron transport chain in photosynthesis as well as in the construction of chloroplasts.

In Chem. Pharm. Bull. 41 (9), 1664-1666, 1993, Shirahashi et al. describe the isolation and identification of anti-tumour-promoting principles from an extract of the cyanobacterium *Phormidium tenue*. Said extract turned out to comprise three classes of glycolipids, that is monogalactosyldiacylglycerol (MGDG), digalactosyldiacylglycerol (DGDG) and sulphoquinovosyldiacylglycerol (SQDG). MGDG and DGDG turned out to inhibit tumour promotion more intensely than SQDG, or rather a mixture of different SQDGs.

The Journal of the National Cancer Institute, Vol. 81, No. 16, August 16, 1989, Gustafson et al., refers to the isolation and identification of four anti-HIV active sulpholipids from cyanobacterial extracts by a combination of gel-permeation and reversed-phase chromatographies. The sulpholipids all had similar levels of activity.


JP patent application No. 64-186626 discloses the use of a sulphonoglycolipid, an SQDG of the formula



to obtain a remedy for intravascular blood coagulation syndrome, and JP patent application No. 64-186627 describes the use of the same compound in the treatment of nephritis.


### Description of the invention

The present invention refers to the use of an SQDG having the formula



wherein  $\text{R}^1$  and  $\text{R}^2$ , which are the same or different, are hydrogen or saturated or unsaturated, optionally hydroxy-substituted acyl groups, protonated or as a salt, for the preparation of a pharmaceutical composition for the treatment of inflammatory skin diseases or disorders.

A preferred SQDG to be used according to the invention is a sulphoquinovosyldiacylglycerol having the formula




wherein  $\text{R}^1$  and  $\text{R}^2$ , which are the same or different, are hydrogen, acetyl or acyl groups comprising 14-22 carbon atoms, preferably 16-18, and having up to 6 unsaturations, preferably 0-3.

As examples of acyl groups  $\text{R}^1$  and  $\text{R}^2$  can be mentioned

fatty acid residues such as palmitoyl, stearoyl, oleoyl, linoleoyl,  $\gamma$ -linolenoyl, columbinoyl, dihomo- $\gamma$ -linolenoyl, arachidon yl, eicosapentaenoyl, docosahexa noyl and ricinoloyl, as well as conjugated forms thereof.

5 The invention specially refers to the use of a 1,2-O,O-dipalmitoyl-[6'-sulpho- $\alpha$ -D-quinovopyranosyl(1'→3)]-sn-glycerol, an SQDG having the formula



15 protonated or as a salt.

Whenever applicable, the expression "an SQDG" is also intended to comprise a salt of SQDG, for instance a sodium, potassium or ammonium salt, or a mixture of one or more SQDG compounds. The counterion can also be a cationic lipid or  
 20 another organic cation to form an SQDG derivative being a prodrug. The SQDG compounds may be isolated from nature as such or may be chemically modified to introduce different acyl groups with specific valuable properties. Examples of modifications are partial or total hydrolysis of acyl groups,  
 25 partial or total hydrogenation, regioselective desaturation and inter- or transesterification. In natural SQDGs the quinovosyl is always in position 3 of the glycerol and has the D-form. The glycosidic link in general has the  $\alpha$ -form.

Especially, the SQDG can be used according to the  
 30 invention for the prophylaxis or treatment of atopic dermatitis, urticaria and psoriasis.

Another use of an SQDG according to the invention is as an antiproliferative agent.

A preferred use of an SQDG according to the invention is  
 35 for the prophylaxis or treatment of psoriasis.

Another preferred use of an SQDG according to the invention, is in combination with a mono- or polyunsaturated fatty acid for the prophylaxis or treatment of psoriasis. In this combination the fatty acids can be in the form of free




fatty acids, monoacylglycerols, diacylglycerols or triacylglycerols, for example a marine oil or evening primrose oil.

5 Mono-unsaturated fatty acids preferably comprise 16-22 carbon atoms, and can for instance be oleic acid.

The polyunsaturated fatty acid, PUFA, to be used in combination with the SQDG should preferably comprise 18-22 carbon atoms and 2-6 unsaturations, and can for instance be an essential fatty acid of the n-6 or n-3 family. Examples of  
10 n-6 acids are  $\gamma$ -linolenic acid, dihomo- $\gamma$ -linolenic acid, arachidonic acid. Examples of n-3 acids are  $\alpha$ -linolenic acid, octadecatetraenoic acid, eicosatetraenoic acid, eicosapentaenoic acid, docosapentaenoic acid and docosahexaenoic acid.

15 In the treatment of atopic dermatitis the fatty acids are preferably chosen among  $\gamma$ -linolenic acid, dihomo gammalinolenic acid and arachidonic acid. In the treatment of psoriasis the fatty acids are preferably chosen among  $\alpha$ -linolenic acid, octadecatetraenoic acid, eicosatetraenoic  
20 acid, eicosapentaenoic acid, docosapentaenoic acid and docosahexaenoic acid.

The invention also refers to a pharmaceutical composition comprising an SQDG having the formula



30

wherein  $\text{R}^1$  and  $\text{R}^2$ , which are the same or different, are hydrogen or saturated or unsaturated, optionally hydroxy-substituted acyl groups, protonated or as a salt, in combination with a polyunsaturated fatty acid and a  
35 therapeutically acceptable carrier for the treatment of dermal diseases, especially psoriasis.

The pharmaceutical composition preferably comprises an SQDG having the formula II or III as described above.


Therapeutically acceptable carriers are for instance

conventional carriers for solutions, suspensions, liposomes, emulsions, aerosols, topical preparations, powders, granules, tablets, capsules or suppositories.

Said pharmaceutical composition can be orally, topically, ocularly, nasally, aurally, vaginally, rectally, enterally or parenterally administered.

In oral administration the administered dose could be from 1 to 100 mg/d, preferably 2-25, especially 2-15 mg/d. In topical application the administered dose should be at least ten times the oral dose giving a preferred range of 10-200 mg/cm<sup>2</sup>/d. The injected amount of the active substance could be estimated to 0.1 to 10 mg/d.

The invention also refers to a new PAF-receptor antagonist, characterized in being an SQDG having the formula



wherein R<sup>1</sup> and R<sup>2</sup>, which are the same or different, are hydrogen or saturated or unsaturated, optionally hydroxy-substituted acyl groups, protonated or as a salt.

SQDG can be obtained from natural sources, such as green plants, plant tissues, sponges, bacteria or algae by drying, such as air drying or lyophilization, extraction and subsequent purification by chromatographic or other methods. In cases where chemical modification of the obtained material is performed, further purification may be necessary to obtain the desired compound. In this application SQDG was obtained by preparing a methanolic extract of the plant material which was subsequently subjected to repeated straight phase column chromatography on silica gel with chloroform and methanol as eluents. Fractions containing polar lipids were then further purified on Sephadex LH20 with water, methanol and ethanol mixtures.

### Description of the drawings

Figure 1 shows the dose response curve of the SQDG of the formula III, and of the PAF receptor antagonist ginkgolide BN 52021;

5        Figure 2 shows the displacement curves of the ginkgolide BN 52021, the PAF receptor antagonist BN 50730, and of the SQDG compound of the formula III.

### Isolation of SQDG from *Polypodium decumanum*

10        The plant material for preparing the Calaguala extract was obtained from a plantation near the Lago de Yohoa in Honduras. *Polypodium decumanum* is a fern which in its natural environment grows as an epiphyte on palm trees, but which is also cultured for its beneficial pharmacological effects.

15        The dried and ground leaves (6.6 kg) were extracted with 3 x 35 l methanol (puriss) overnight under stirring at room temperature. The extract was filtered, concentrated in vacuo and lyophilised (yield 267 g). 50 g of the methanolic extract was then dissolved in 400 ml water and extracted with 4 x 600  
20        ml chloroform; yield 16 g. The chloroform extract was adsorbed on silica gel (Merck, mesh 70-230) and eluted through a column packed with 200 g silica gel using a gradient of chloroform to methanol. Fractions of 20 ml were collected and pooled together according to TLC comparison. 10  
25        fractions were obtained and checked for activity in the elastase assay described below. The most active fractions (1.8 g) were repeatedly chromatographed on silica gel with a gradient of chloroform to methanol and the combined active fractions (1.4 g) were then eluted through a Sephadex LH 20  
30        column (Pharmacia Fine Chemicals, Sweden) with a gradient of methanol (30%) and water to 100 % methanol as mobile phase. The next to the last fraction (210 mg) showed the highest activity in the assay and was further purified using a second  
35        Sephadex LH 20 column which was first washed with ethanol and then eluted with methanol. By this 10 mg of the compound 1,2-O,O-dipalmitoyl-[6'-sulpho- $\alpha$ -D-quinovopyranosyl(1'→3)]-sn-glycerol, that is a compound of the formula III above, was obtained. The structure of the compound was verified as below.

# Identification of the SQDG compound

The structure of the compound was identified by means of mass spectrometry, carbohydrate analysis and NMR.

GC-MS analysis was done on a HP 5890 gas chromatograph coupled to a HP 5970A quadrupole mass selective detector operated at an ionization voltage of 70 eV and an electron multiplier voltage of 1800 V. The compound was analyzed after a transesterification reaction according to Seppänen-Laakso, T., et al., Acta Pharm. Fenn. 99, 109-117, 1990, which allows the identification of bound fatty acids. A single peak was observed in the GC chromatogram and the peak exhibited major signals at  $m/z$  ( $M^+$ ) 239, 227, 213, 199, 185, 157, 143, 129, 97, 87, 74, 59, 43 and was thus identified as the methylester of palmitic acid. For the molecular weight determination a positive ion FAB mass spectrum was recorded with a JEOL SX 102 instrument with glycerol as matrix. By this a molecular ion  $[M+1]^+$  at  $m/z = 817$  was displayed corresponding to a molecular formula of  $C_{41}H_{77}O_{12}SNa$  for the compound.

Carbohydrate analysis was performed by hydrolysing the compound with trifluoroacetic acid followed by reduction and acetylation and the resulting peracetylated compound was compared by gas chromatography with authentic carbohydrates treated in the same manner. This comparison yielded glucose.

The NMR experiments were run using a JEOL 270 MHz instrument and a mixture of  $CD_3OD$  and  $CDCl_3$  as solvent and TMS as internal standard. The  $^1H$ -NMR and  $^{13}C$ -NMR shift values for the compound are shown in Table 1. The values have been arranged into three groups, that is acyl, glycerol and sulphoquinovosyl moieties.

Table 1. NMR-values

	<u><math>^1H</math> shift</u>	<u><math>^{13}C</math> shift</u>
Acyl moieties		
C16	0.88(t)	14.3
C15		23.4
C14		32.7
C methylene	1.25(m)	29.9, 30.2, 30.5
C3	1.59(m)	25.7
C2	2.3(b)	34.8, 35.0
C1		173.7

## Glycerol moiety

sn-1*	4.49(dd), 4.17(dd)	64.0
sn-2	5.3(m)	71.3
sn-3	4.08(dd), 3.57(dd)	66.7

5

## Sulphoquinovosyl moiety

C1'	4.81(d)	99.0
C2'	3.40(dd)	73.0
C3'	3.62(t)	74.6
C4'	3.10(t)	74.6
C5'	4.05(m)	69.4
C6'	3.34(dd), 2.94(dd)	54.0

10

\* sn refers to stereospecific numbering

15

## Formulations

Oral emulsion

An oil-in-water emulsion (batch size 250 g) was prepared with the following ingredients:

20

<u>Ingredients</u>	<u>%</u>
Corn oil	20.00
CPL <sup>®</sup> -Galactolipid	1.70
SQDG	0.15
25 Sucrose	17.00
Potassium sorbate	0.20
Ascorbyl palmitate	0.02
Ammonium phosphatides	0.10
Lemon flavour	0.20
30 Water	ad 100.00

The emulsifier (CPL<sup>®</sup>-Galactolipid, a lipid emulsifier manufactured by Scotia LipidTeknik AB, Sweden) and antioxidant were dispersed in the oil. Sucrose, preservative, flavour and water were mixed. The oil phase and the aqueous phase were preheated to 50°C and then the oil was added to the aqueous phase under high shear mixing at 16,000 rpm for 3.5 min. The preemulsion was then homogenised at 80 MPa and 55°C for 7 cycles (Mini-Lab 8.30 H; APV Rannie AS, Denmark).

40 This formulation resulted in a milky emulsion. The daily dose of this emulsion is 10 ml.

The neutral vegetable corn oil can be replaced by for instance an oil rich in  $\gamma$ -linolenic acid, such as evening primrose oil, EPO, giving an emulsion comprising SQDG and PUFA.

5

#### Topical formulation

	<u>Ingredient</u>	<u>g</u>
	CPL <sup>®</sup> -Galactolipid	17.00
	SQDG	2.00
10	Metagin	0.05
	Propagin	0.01
	Water	ad 100.00

15 SQDG was dispersed in water containing the preservatives. After the addition of the galactolipids, prepared from oats, the mixture was alternatively vortexed and stirred giving a viscous dispersion apted for topical application.

#### **Biological tests**

20 In vivo animal models to test for anti-psoriatic activity are few and inadequate. Therefore, models that focus on the different characteristics of the disease are frequently used. One possibility is to consider the reported immunological background of the disease and the fact that the  
25 immunosuppressive drug cyclosporin A shows excellent therapeutic effects in treatment of severe psoriasis to use a test model where the ability of a compound to suppress the immune system is studied. In Phytother. Res. 5, 234-236, 1991, Tuominen et al. report an enhancing effect of Calaguala  
30 on the prevention of rejection of skin transplants in mice.

The ability of Calaguala methanolic extract to retard rejection of skin grafts in mice was studied using an adaptation of a method described by Billingham et al., J. Exp. Biol. 28, 385-403, 1957. Mice were pre-treated with  
35 either Calaguala orally or subcutaneously or with cyclosporin A subcutaneously for two weeks whereafter they were grafted with skin from a donor mice of different strain. They were then examined daily for signs of rejection. The results show that the Calaguala methanolic extract administered in a dose

of 500 mg/kg/day gave a significant enhancement of the rejection time as compared to untreated animals. The mechanism for this effect could be the same as for the clinical effects reported for treatment of psoriasis with Calaguala.

The inflammatory aspect of psoriasis can be studied by using arachidonic acid induced oedema in rat ear, Bosman, Skin. Pharmacol. 7, 324-334, 1994, and by using the rat paw oedema test, a classical way of experimentally investigating the effect of different compounds in acute inflammation.

The inflammatory tests described below were performed with the methanolic extract obtained from Polypodium decumanum, as described above, the Calaguala extract, in order to evaluate an optional activity of the extract on psoriasis.

It has previously been shown that the Calaguala extract exhibits dose-dependent inhibition of leukotriene B<sub>4</sub> synthesis in human leucocytes. This inhibitory activity was found to be caused by the polyunsaturated fatty acids linoleic, linolenic and arachidonic acid. The activity of additional unsaturated fatty acids, such as oleic acid and eicosapentaenoic acid, was also examined in the model and were all found to exhibit comparable inhibitory patterns. Leukotriene B<sub>4</sub> is an inflammatory mediator isolated in abnormally high quantities in the psoriatic skin. See Vasänge et al., Prostaglandins, Leukotrienes and Essential Fatty Acids 50, 279-284, 1994.

#### Rat ear oedema

The rat ear oedema test is aimed to be used for detection of drugs with topical anti-inflammatory activity. It is based on the work by Tonelli et al., Endocrinology 77, 625-634, 1965, but instead of using the carcinogenic croton oil as induced, ethyl phenyl propiolate (EPP) is applied topically on the rat ears. The oedema develops rapidly (about 30 minutes) and starts to decline after two hours. Several measurements can be made by using a special measuring device and the values obtained with application of both EPP and the drug to be investigated are compared to values obtained with

EPP alone.

The mechanism of the EPP induced oedema is not completely understood but it is known that compounds with broad anti-inflammatory action, like steroidal anti-inflammatory drugs, are active in this assay. The oedema can also be induced by other agents, such as arachidonic acid, and in that specific case the oedema is believed to develop through the action of the phospholipid, inflammatory mediators, leukotrienes. They can also contribute to the EPP oedema.

The Calaguala methanolic extract was tested in this model on several occasions. It was applied either locally (dissolved in methanol or acetone) or orally. Representative experiments are presented below:

**Table 2. Ear thickness**

<u>Treatment</u>	<u>Number of ears treated</u>	<u>Mean decrease in oedema</u>
None	6	0
Calaguala 400 µg/ear	12	30%
Mepyramin	6	44%

The thickness of ears is measured after 2 hours. As positive control the antihistamine mepyramine (1 µg/ear) is used. The significance of the values obtained was determined by using Students t-test (one tailed analysis). For mepyramin the significance was  $p < 0.01$  and for Calaguala  $p < 0.001$ .

**Table 3. Ear thickness**

<u>Treatment</u>	<u>Number of ears treated</u>	<u>Mean decrease in oedema</u>
None	20	0
Topical Calaguala	20	32%
Pre-treatment Calaguala	20	32%

The thickness of the ears was measured after 2 hours. One Calaguala group received the extract topically, 400 µg/ear, and another group had been pre-treated for two weeks with about 1000 mg/kg/d Calaguala in their drinking water. The significance was  $p < 0.001$  for the local treatment and  $p < 0.005$  for the oral treatment (Students t-test, one tailed).



Rat paw oedema

The rat paw oedema test is a classical way of experimentally investigating the effect of different compounds in acute inflammation. The most commonly used inducer is carrageenan which is injected to the hind paw of male Sprague Dawley rats. The mechanism of the oedema development which culminates after three hours is not completely understood. In general, it is believed to involve stimulation of prostaglandin synthesis, a fact that is supported by the inhibitory activity of cyclo-oxygenase inhibitors in this assay. More recent data, however, indicate that additional mechanisms are involved in the oedema development. For example, it has been shown that early phase of the carrageenan induced rat paw oedema is partly dependent on PAF release (Caruso et al., Pharmacol. Res. 31(1), 67-72, 1995). PAF has also been found to potentiate the oedema induced by carrageenan and the known PAF receptor antagonist, BN 52021, is capable of abolishing this enhancement.

Several tests were performed with the methanolic Calaguala extract which was applied orally in doses 500-2000 mg/kg. As positive control the cyclo-oxygenase inhibitor Na-salicylate (200 mg/kg) was used. A representative study is presented below:

Table 4. Effect of Calaguala methanolic extract on rat paw oedema induced by carrageenan.

Treatment	No. of animals	Mean decrease in oedema	Significance
-----------	----------------	-------------------------	--------------

None	5	-	-
Na-salicylate	5	43%	p<0.0005
Calaguala 500 mg/kg	5	19%	p<0.05
Calaguala 1000 mg/kg	5	22%	p<0.025
Calaguala 2000 mg/kg	5	21%	p<0.05

The significanci s were calculated using Students t-test (one tailed).

In another set of experiments rats were pre-treated for

two weeks with Calaguala methanolic extract in their drinking water. No statistically significant decrease in the oedema formation was observed when the rats were challenged with carrangeenan on day 14. This finding is well in line with the theory of PAF being involved in the early phase of the carrageenan induced oedema whereas the later phases would be caused by stimulation of prostaglandins.

The Calaguala extract has also been investigated for inhibitory activity on the enzyme cyclooxygenase using bovine seminal vessel microsomes (White et al., Prostaglandins 7, 123-9, 1974). Cyclooxygenase is the first enzyme in the reaction chain for conversion of arachidonic acid to prostaglandins. The extract failed to show any inhibitory activity in this assay. This could be the explanation to why the inhibition caused by Calaguala in the carrageenan induced rat paw oedema does not increase with increased concentrations of the extract: Only a minor part of the oedema increase is caused by PAF and Calaguala is only capable of inhibiting that part leaving the increase prostaglandin production unaffected. Thus the concentration of 500 mg/kg is already sufficient to cause total inhibition of the PAF induced part.

#### Quantification of SQDG in Calaguala methanolic extract

Methanolic extract prepared from leaves of Polypodium decumanum has been analysed for its content of sulphonoglycolipid, SQDG.

5700 gram of finely ground plant material was extracted with methanol overnight, under stirring at room temperature. The process was repeated three times with a total of 35 liter methanol. The extract was filtered and dried and the yield was 337 gram, In order to get a detectable signal for SQDG the methanolic extract was subjected to a preliminary fractionation.

Initially the extract was dissolved in water and the lipids were separated from the polar components by partitioning to chloroform. The chloroform extract was thereafter applied on a Sephadex LH20 column which had been equilibrated with ethanol. The column was first eluted with

ethanol and thereafter with methanol. SQDG was selectively enriched in the methanol fraction. The fractions eluting prior to the change of eluent was examined by NMR for the presence of SQDG but only minor traces could be detected.

5       The HPLC analysis was carried out by using a straight phase system with polyvinyl alcohol bonded silica column (T-column, 89°C, PVA-SIL, 250x4.6 mm, 5µm), a flow rate of 1 ml/min, a light scattering detector (Sedex 45) and a mobile phase with a gradient system with hexane, iso-propanol, n-  
10 butanol, tetrahydrofuran, iso-octane, water and ammonium acetate.

Commercially available SQDG was used as external standard and regression analysis was used to obtain an equation for a straight line. This was then used to calculate the amounts of  
15 SQDG present in the extract.

The amount SQDG present in the Calaguala extract was determined to be 4 mg/g extract.

#### PAF induced exocytosis.

20       In this test, which is also called the elastase assay, the effect of the extracts on inhibiting the exocytosis induced by PAF was examined. This model was used to guide the fractionation of the extracts or isolated compounds.

Cell preparation: Peripheral blood anti-coagulated with  
25 heparin was obtained from healthy volunteers at the University Hospital (in Uppsala, Sweden). After sedimentation with 10 % Dextran T-500 (Pharmacia Fine Chemicals) for 30 min at 20°C the supernatant was removed and centrifuged at 200 x g for 30 min. The sediments were treated with one volume of  
30 ice-cold water for 21 s followed by the addition of 9 volumes of  $Mg^{2+}$  and  $Ca^{2+}$  free PBS in order to lyse the remaining erythrocytes. After centrifugation (200 x g for 10 min at 4°C) the leucocytes were suspended in PBS (containing  $Mg^{2+}$  and  $Ca^{2+}$  and Cytochalasin B, Serva, 5 µg/ml) at a  
35 concentration of  $10-30 \times 10^6$  cells/ml.

The assay was performed as described by Tuominen et al., Planta Medica 58, 306-310 (1992). In short leucocytes were incubated at 37°C in PBS containing 2.5 % BSA together with SAAVNA (N-succinyl-L-alanyl-L-alanyl-L-valine-p-nitroanilide,

Bachem), the inhibitor in different concentrations or the vehicle (max. 0.1 % DMSO) and PAF (Bachem) for 10 min. Blank tubes without PAF were run in parallel. The reaction was stopped by the addition of 2 % citric acid and after the centrifugation, the samples were measured in a UV spectrophotometer at 405 nm. The absorbance of the corresponding blank tube was subtracted from the sample and the inhibition of PAF induced elastase release was calculated as the relative increase in absorbance as compared to the vehicle. The samples were analyzed as 4-5 concentrations ranging from  $10^{-3}$  M to  $10^{-7}$  M. They were dissolved in 10 % DMSO (the isolated compound) or cyclodextrin (BN 52021, used as a reference) and then diluted with the buffer. The final concentration of DMSO or cyclodextrin never exceeded 0.1 %.

The dose response curve of the SQDG of the formula III, and of the PAF receptor antagonist ginkgolide BN 52021 is shown in Figure 1. Each point represents the mean of 2-4 experiments, all performed in duplicate. The  $IC_{50}$  value was determined to 10  $\mu$ M for the compound of the invention and to 80  $\mu$ M for BN 52021.

In another experiment, the activity of SQDG rich fractions of different origin, such as from Spirulina and from bladder-wrack, were compared to the activity of SQDG from Calaguala, but no differences could be observed.

In still another experiment different polar lipids were tested in the model. Compounds such as MGDG and DGDG showed no inhibitory activity in concentrations around the  $IC_{50}$  of SQDG.

### [ $^3$ H]PAF binding assay

The PAF induced exocytosis model described above is used for screening for PAF receptor antagonists since it is a rapid, reproducible and reliable so called functional assay for PAF. However, a number of other actions, like elastase inhibition or membrane stabilisation, would also result in an inhibitory effect in the model which consequently is not exclusive for PAF receptor antagonism. To elucidate if SQDG acts through PAF receptor antagonism in the neutrophil, a receptor binding model using [ $^3$ H]PAF as the radioligand was

used.

Neutrophils are the most abundant circulating white blood cells and are usually the first cells to respond to an infectious or inflammatory stimulus. They are capable of phagocytosing appropriately opsonized bacteria and may also upon stimulation with e.g. PAF, secrete potentially toxic oxygen metabolites and lysosomal enzymes, including elastase, into the supernatant medium. These cells can also adhere to vascular endothelium and migrate towards the source of the chemoattractants. All these properties are a part of the response of the organism to inflammatory or infectious disturbance.

Cell preparation: For the receptor binding experiments the cells were prepared in the same way as described above except that concentrated suspensions of human leukocytes in a CDP-adenine solution were used and in the final step the cells were suspended in the incubation buffer and the cell count was adjusted to  $3.1 \times 10^6$  cells/ml.

[ $^3\text{H}$ ]PAF ( $\text{C}_{18}$ ; 1-O-[ $^3\text{H}$ ]octadecyl-2-acetyl-sn-3-phosphocholine; Amersham, England) receptor binding was studied in human polymorphonuclear leukocytes ( $2.5 \times 10^6$  cells/ml) using a method recently reevaluated by us (Vasänge et al., in press, 1996). Incubations were performed for 120 min at  $20^\circ\text{C}$  in a total volume of 1.0 ml containing 0.6 nM  $\text{Na}_2\text{HPO}_4$ , 0.6 nM  $\text{NaH}_2\text{HPO}_4$ , 25 mM Tris HCl, 130 mM NaCl, 5.5 mM KCl, 1.4 mM  $\text{CaCl}_2$ , 0.7 mM  $\text{MgCl}_2$ , 10 mM glucose, 0.5 % BSA and 0.35-0.40 nM [ $^3\text{H}$ ]PAF. Final pH was 7.0 at the incubation temperature used. Non-specific binding was determined with 1  $\mu\text{M}$  C16-PAF (Bachem, Switzerland). The incubation contents were rapidly filtered by vacuum over Whatman GF/C glass-fibre filters using a 12 well Skatron Cell Harvester (Skatron A/S, Norway) and washed with 3 x 2 ml incubation buffer containing 0.1 % BSA. The filtration took less than 15 s. Filters were also pre-soaked with incubation buffer containing 0.1 % BSA. Following filtration, the filters were dried in an oven at  $70^\circ\text{C}$  for 30 min and equilibrated for two hours in the scintillation vials before counting in a Packard scintillator at an efficiency of 55 %. The competition binding assays were performed in triplicate in several concentrations ( $10^{-11}$  -

10<sup>-3</sup> M) of the isolated compound and the receptor antagonists BN 50730 and BN 52021 as references. The compound and BN 50730 were dissolved in 10 % DMSO (10<sup>-2</sup> M), BN 52021 in cyclodextrin and then further diluted with the incubation  
5 buffer.


The displacement curves of the ginkgolide BN 52021, the hetrapazine-derived PAF receptor antagonist BN 50730 and of the SQDG compound of the formula III are presented in Figure 2. The figure shows the decrease on dpm measured as  
10 percentage of total bound ligand (Bt). The level of non-specific binding was determined to about 45 % of the Bt. Each point represents the mean of 2-4 experiments, all performed in triplicate. The IC<sub>50</sub> value was determined to be 2 µM for the isolated compound, 25 µM for BN 52021 and 0.04 µM for BN  
15 50730.

#### Summary

Calaguala has in clinical trials been shown to have beneficial effects in the treatment of psoriasis. One of the  
20 pathological findings in the psoriatic skin is elevated amounts of the inflammatory mediator platelet activating factor. The crude methanolic extract of Calaguala which gave the effects in psoriasis also shows a dose dependent inhibition of the PAF induced elastase release in human  
25 neutrophils, which is a functional assay for PAF receptor antagonism. When activity guided fractionation of the Calaguala extract was carried out, the main compound responsible for the PAF activity was discovered to be SQDG. SQDG was also proven to act through binding to PAF receptors  
30 in human neutrophils indicating a true receptor antagonistic property. Thus SQDG is believed to have therapeutic potential in treatment of diseases, such as inflammatory skin diseases. Another pathological finding in the psoriatic skin is elevated amounts of leukotriene B<sub>4</sub> which is known to be  
35 inhibited by polyunsaturated fatty acids. The combination of unsaturated fatty acids and SQDG, as in the Calaguala extract, should therefore be beneficial in the treatment of skin inflammation.


## CLAIMS

1. Use of an SQDG having the formula




wherein  $\text{R}^1$  and  $\text{R}^2$ , which are the same or different, are hydrogen or saturated or unsaturated, optionally hydroxy-substituted acyl groups, protonated or as a salt, for the preparation of a pharmaceutical composition for the treatment of inflammatory skin diseases or disorders.

2. Use according to claim 1 of a sulphoquinovosyl-diacylglycerol, an SQDG having the formula



wherein  $\text{R}^1$  and  $\text{R}^2$ , which are the same or different, are hydrogen, acetyl or acyl groups comprising 14-22 carbon atoms, preferably 16-18, and having up to 6 unsaturations, preferably 0-3.

3. Use according to claim 1 or 2 of a 1,2-O,O-dipalmitoyl-[6'-sulpho- $\alpha$ -D-quinovopyranosyl(1'→3)]-sn-glycerol, an SQDG having the formula



protonated or as a salt.

4. Use according to any of claims 1-3 of an SQDG for the prophylaxis or treatment of atopic dermatitis, urticaria or psoriasis.


5. Use according to any of claims 1-3 of an SQDG as an antiproliferative agent.

6. Use according to any of claims 1-3 of an SQDG for the prophylaxis or treatment of psoriasis.

7. Use according to any of claims 1-3 of an SQDG in combination with a polyunsaturated fatty acid for the prophylaxis or treatment of psoriasis.

8. Use according to claim 7 of an SQDG in combination with linoleic acid or linolenic acid.


9. Pharmaceutical composition comprising an SQDG having the formula



wherein  $\text{R}^1$  and  $\text{R}^2$ , which are the same or different, are hydrogen or saturated or unsaturated, optionally hydroxy-substituted acyl groups, protonated or as a salt, in combination with a polyunsaturated fatty acid and a therapeutically acceptable carrier.

10. Pharmaceutical composition according to claim 9, comprising an SQDG having the formula






5

wherein  $\text{R}^1$  and  $\text{R}^2$ , which are the same or different, are hydrogen, acetyl or acyl groups comprising 14-22 carbon atoms, preferably 16-18, and having up to 6 unsaturations, preferably 0-3, and a polyunsaturated fatty acid from the n-3 family for the treatment of psoriasis.

10

11. Pharmaceutical composition according to claim 9, comprising an SQDG having the formula

15




20

wherein  $\text{R}^1$  and  $\text{R}^2$ , which are the same or different, are hydrogen, acetyl or acyl groups comprising 14-22 carbon atoms, preferably 16-18, and having up to 6 unsaturations, preferably 0-3, and a polyunsaturated fatty acid from the n-6 family for the treatment of atopic dermatitis.

25

12. Pharmaceutical composition according to any of claims 9-11, comprising a 1,2-O,O-dipalmitoyl-[6'-sulpho- $\alpha$ -D-quinovopyranosyl(1'→3)]-sn-glycerol, an SQDG having the formula

30




35

protonated or as a salt.

13. Pharmaceutical composition according to any of claims 9-12 for oral, topical, rectal, enteral or parenteral administration.

- 5 14. A new PAF-receptor antagonist, characterized in being an SQDG having the formula



- 15 wherein  $\text{R}^1$  and  $\text{R}^2$ , which are the same or different, are hydrogen or saturated or unsaturated, optionally hydroxy-substituted acyl groups, protonated or as a salt.

20

25

30

35

1/2




FIG. 1




FIG. 2

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 97/00735

<b>A. CLASSIFICATION OF SUBJECT MATTER</b>		
IPC6: A61K 31/70 According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b>		
Minimum documentation searched (classification system followed by classification symbols)		
IPC6: A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
SE,DK,FI,NO classes as above		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
CA, WPI, MEDLINE		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	J. Pharm. Pharmacol., Volume 49, 1997, Mervi Vasänge et al, "A Sulphonoglycolipid from the Fern Polypodium decumanum and its Effect on the Platelet Activating-factor Receptor in Human Neutrophils" page 562 - page 566  --	1-12
A	Chem. Pharm. Bull., Volume 41, No 9, 1993, Takashi Morimoto et al, "Studies on Glycolipids. VII. Isolation of Two New Sulfoquinovosyl Diacylglycerols from the Green Alga Chlorella vulgaris" page 1545 - page 1548	1-8
X	--	9-14
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
5 August 1997		28 -08- 1997
Name and mailing address of the ISA/ Swedish Patent Office Box 5055, S-102 42 STOCKHOLM Facsimile No. +46 8 666 02 86		Authorized officer  Eva Johansson Telephone No. +46 8 782 25 00

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 97/00735

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Journal of the National Cancer Institute, Volume 81, No 16, August 1989, Kirk R. Gustafson et al, "AIDS-Antiviral Sulfolipids From Cyanobacteria (Blue-Green Algae)" page 1254 - page 1258	9-14
A	--	1-8
X	Patent Abstracts of Japan,, abstract of JP,A, 3-52815 (EISAI CO LTD), 7 March 1991 (07.03.91)	9-13
A	--	1-8,14
X	Patent Abstracts of Japan,, abstract of JP,A, 3-52816 (EISAI CO LTD), 7 March 1991 (07.03.91)	9-13
A	-----	1-8,14